

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/50, A61K 38/48	A1	(11) International Publication Number: WO 98/38291 (43) International Publication Date: 3 September 1998 (03.09.98)												
(21) International Application Number: PCT/GB98/00590 (22) International Filing Date: 25 February 1998 (25.02.98) (30) Priority Data: <table border="0"><tr><td>9703850.9</td><td>25 February 1997 (25.02.97)</td><td>GB</td></tr><tr><td>9703827.7</td><td>25 February 1997 (25.02.97)</td><td>GB</td></tr><tr><td>9704252.7</td><td>28 February 1997 (28.02.97)</td><td>GB</td></tr><tr><td>9706119.6</td><td>25 March 1997 (25.03.97)</td><td>GB</td></tr></table> (71) Applicant (for all designated States except US): CORTECS (UK) LIMITED [GB/GB]; The Old Blue School, Lower Square, Isleworth, Middlesex TW7 6RL (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): MYNOTT, Tracey, Lehanne [AU/GB]; Imperial College of Science, Technology & Medicine, Dept. of Biochemistry, Exhibition Road, London SW7 2AZ (GB). ENGWERDA, Christian [AU/GB]; London School of Hygiene & Tropical Medicine, Dept. of Medical Parasitology, Keppel Street, London WC1E 7HT (GB). PEEK, Keith [GB/GB]; Cortecs (UK) Limited, Research & Development Division, Techbase 1, Newtech Square, Deeside Industrial Park, Deeside, Clwyd CH5 2NT (GB).		9703850.9	25 February 1997 (25.02.97)	GB	9703827.7	25 February 1997 (25.02.97)	GB	9704252.7	28 February 1997 (28.02.97)	GB	9706119.6	25 March 1997 (25.03.97)	GB	(74) Agents: CHAPMAN, Paul, William et al.; Kilburn & Strode, 20 Red Lion Street, London WC1R 4PJ (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
9703850.9	25 February 1997 (25.02.97)	GB												
9703827.7	25 February 1997 (25.02.97)	GB												
9704252.7	28 February 1997 (28.02.97)	GB												
9706119.6	25 March 1997 (25.03.97)	GB												
(54) Title: COMPONENT OF BROMELAIN (57) Abstract The invention relates to a component of bromelain which is largely responsible for the ability of bromelain to interrupt the MAP kinase cascade. The component contains ananain and comosain and is useful in the treatment or prevention of diseases and conditions mediated by T cell activation or by activation of the MAP kinase pathway.														

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

COMPONENT OF BROMELAIN

5 The present invention relates to a component of bromelain. In particular, the invention relates to the use of this bromelain component in medicine, particularly as an anti-cancer agent and an immunosuppressive agent.

10 Stem bromelain (bromelain) is the collective name for the proteolytic enzymes found in the tissues of the plant *Bromeliaceae*. It is a mixture of various moieties derived from the stem of the pineapple plant (*Ananas comosus*). Bromelain is known to contain at least five proteolytic enzymes but also non-proteolytic enzymes, including an acid phosphatase and a peroxidase; it may also contain amylase and cellulase activity. In addition, various other components are present.

15 Bromelain has previously been used in the treatment of a variety of conditions including inflammation and, in particular, it has been used in the treatment of diarrhoea. The use of bromelain in the treatment of infectious diarrhoea is described in WO-A-9301800, where it is suggested that bromelain works by destroying intestinal receptors for pathogens by proteolysis, and in WO-A-8801506, which teaches that bromelain detaches pathogens from intestinal receptors.

20 Taussig *et al*, *Planta Medica*, 1985, 538-539 and Maurer *et al*, *Planta Medica*, 1988, 377-381 both suggest that bromelain may be of use in inhibiting tumour growth. US 5,223,406, DE-A-4302060 and JP-A-59225122 also teach the use of bromelain in the treatment of cancer. US 5,223,406 teaches that bromelain is capable of inducing
25 tumour necrosis factor (TNF) while DE-A-4302060 teaches that bromelain can prevent metastasis by the structural modification of the tumour surface protein CD44.

In WO-A-9400147, various experiments were described which demonstrate that

proteolytic enzymes and, in particular, bromelain, are capable of inhibiting secretion. The application also discloses that bromelain can reduce toxin binding activity and can inhibit the secretory effect of toxins such as heat labile toxin (LT) and cholera toxin (CT) and also toxins such as heat stable toxin (ST). This is in spite of the fact that ST has a very different mode of action from LT and CT. These observations were explained by the fact that one component of the bromelain mixture, stem bromelain protease, appears to be capable of modulating cyclic nucleotide pathways and this is discussed further in WO-A-9500169. In addition, bromelain has also been demonstrated to inhibit secretion caused by the calcium dependent pathway.

The present inventors have studied the varied biological effects of bromelain and, in particular, its effects in a well documented model of intracellular signal transduction, namely T cell receptor (TCR)/CD3 signalling and IL-2 production. Significant progress over recent years has led to the understanding of biochemical events which occur following TCR engagement (reviewed Cantrell, *Annu. Rev. Immunol.* **14**, 259-274, (1996)), therefore TCR signalling provides an excellent model for elucidation of the effects of biologically active compounds. Effective T cell activation requires two signals. The first signal is generated by the TCR/CD3 complex after engagement with antigen peptide presented by the major histocompatibility complex (MHC) expressed on antigen presenting cells (APC) (Cantrell, 1996). The second, costimulatory signal is generated by ligation of CD28 receptors on T cells with the B7 family of ligands on APC. A key element in the signalling pathway involved in transducing receptor-initiated signals to the nucleus is the family of mitogen-activated protein kinases (MAPk). The best studied of these kinases are the extracellular signal-regulated protein kinases (ERK)-1 and ERK-2 (also referred to as p44^{MAPk} and p42^{MAPk}, respectively). ERK's are serine/threonine kinases that are activated when phosphorylated on tyrosine and threonine residues. *In vitro*, this activation is reversed if either residue is dephosphorylated. A relatively newly discovered member of the MAPk family are c-

Jun NH₂-terminal kinases (JNKs) which exist as 46 kDa and 55 kDa forms that also require phosphorylation for activation. ERK activation is dependent on p56^{Lck} and coupling of the TCR/CD3 complex to p21^{Ras}, with subsequent activation of the Raf-1/MEK1/ERK kinase cascade. JNK activation also requires p21^{Ras}, as well as signals
5 generated by the CD28 costimulatory receptor which activate GTP (guanosine triphosphate)-binding proteins (such as Rac1 or Cdc42) that induce the PAK/MEKK/SEK/JNK kinase cascade. Activated ERK phosphorylates Elk-1, which in turn, mediates induction of c-fos activity following phosphorylation of c-jun by JNK. Activated c-fos and c-jun combine to form the AP-1 protein required for IL-2
10 synthesis. The above events are summarised in Figure 1. All the above-mentioned signalling events require tyrosine phosphorylation, as inhibitors of protein tyrosine kinases (PTKs) inhibit many events associated with TCR stimulation, including T cell activation and IL-2 production.

15 In WO-A-9600082, we showed that bromelain could inhibit tyrosine phosphorylation and activation of ERK-2 in T cells stimulated via the TCR, or with combined phorbol ester plus calcium ionophore. We have now found that, in association with decreased ERK activity, bromelain decreased IL-2, IL-4 and IFN- γ mRNA accumulation in T cells stimulated with phorbol ester and ionophore, but did not affect cytokine mRNA
20 accumulation in cells stimulated via the TCR. This data suggests the existence of a TCR-activated, ERK-independent pathway involved in cytokine production in T cells.

From the prior art, it is clear that bromelain is a mixture which has a variety of different physiological effects. Not all of the components of the bromelain mixture
25 have been characterised and so, except for stem bromelain protease, whose activity we have described, it is not clear which of the components is responsible for which of the various different effects of bromelain. This is, of course, a major disadvantage if the bromelain mixture is to be administered as a pharmaceutical because while one

component of bromelain might give the desired effect, there may well be unwanted side effects arising from the action of some other component of the bromelain mixture.

5 It would therefore be beneficial if individual components of bromelain giving rise to particular medicinal activities could be isolated and administered separately so as to lessen the possibility of side effects. We have now identified an active fraction of crude bromelain which is responsible for its ability to inhibit ERK activation, and therefore block the MAP kinase pathway. Although not a single protein, this fraction consists of only a few components and so the possibility of side effects when it is
10 administered to patients is greatly reduced compared with crude bromelain.

The fraction of the invention, which the inventors have designated CCS, may be isolated from the bromelain mixture by conventional methods, for example by chromatography. High performance liquid chromatography (HPLC) is suitable for the
15 purpose and particularly good separation of bromelain proteins may be achieved by fast protein liquid chromatography (FPLC™) using a column packing material such as S-sepharose. As will be described in more detail in the examples, in chromatography on S-sepharose using a linear gradient of 0 to 0.8M sodium chloride in acetate buffer over 300ml, the protein of the present invention was the last double peak off the column.

20

In a first aspect of the present invention, there is provided a component of bromelain which contains proteins having molecular weights of about 15.07kDa, 25.85kDa and 27.45kDa as determined by SDS-PAGE, has isoelectric points of 10.4 and 10.45 and is obtainable by the following method:

25

- i. dissolving bromelain in acetate buffer at pH 5.0;
- ii. separating the components of the bromelain by fast flow high

performance liquid chromatography on S-sepharose eluting with a linear gradient of 0 to 0.8 M sodium chloride in acetate buffer over 300 ml;

iii. collecting the fraction corresponding to the final double peak off the column; and

iv. isolating the protein from the fraction collected in (iii).

This fraction, termed CCS, has been found to have a number of potentially useful activities. Firstly, we have found that it blocks ERK-2 phosphorylation, and therefore the MAP kinase cascade. In addition, it blocked IL-2 production and CD4⁺ T cell proliferation. However, CCS did not affect splenocyte proliferation, which suggests that it has a selective mode of action. CCS also differentially blocked growth of human tumour cell lines including ovarian, lung, colon, melanoma and breast tumours. The differential activity of CCS against the different cell lines further suggests that CCS has a selective mode of action and that it may also act as an anti-cancer agent. The inhibitory effect of CCS on ERK-2 is dependent on its proteolytic activity, since E-64, a selective cysteine protease inhibitor, could abrogate the effect of CCS.

Although in WO-A-9724138, we stated that proteases in general are capable of decreasing MAP kinase activation, we have now found that this is not the case as trypsin does not abrogate T cell signalling and, indeed, in other studies, has been shown to increase MAPk activation (Belham *et al*, 1996, *Biochem. J.*, **320**, 939-946). Thrombin, a protease involved in the blood coagulation cascade, has also been shown to increase MAP kinase activation (Vouret-Craviari *et al*, 1993, *Biochem. J.*, **289**, 209-214). The inventors have now also shown that other proteases contained within the crude bromelain mixture do not block the activation of the MAP kinase pathway.

It is possible that the effects of CCS on the MAP kinase pathway in T cells are mediated by specific proteolytic effects at the cell surface. It is known that bromelain cleaves the CD45 RA isoform and selectively removes other surface molecules from human PBMCs. Bromelain also partially removes CD4 from T cell surfaces. Since
5 CD45 and CD4 play an obligate stimulatory role in TCR-mediated T cell activation, CCS may interfere with TCR signalling by affecting these molecules. Although the importance of CD45 and CD4 is well recognised for TCR-initiated signal transduction, it is possible to bypass their requirements for T cell activation by the use of phorbol ester and calcium ionophore. Use of combined phorbol esters and
10 ionophore restores normal function to T cells which have been made refractory to TCR stimulation by the use of tyrosine kinase inhibitors or which are CD45 or p56^{Lck} deficient.

However, in the present study, the inventors have shown that normal function is not
15 restored to T cells pre-treated with CCS when they are treated with PMA plus ionophore. The inhibitory effect of CCS on ERK-2 is thus not thought to be mediated *via* effects on CD45 or CD4 on T cells. CCS possibly affects an as yet unidentified surface molecule, which, in turn, affects the MAP kinase pathway. The inhibitory effect of CCS on cytokine production is thus not thought to be mediated
20 *via* its effects on CD45 or CD4 on T cells. The inhibitory effect of CCS on T cell signal transduction was not because of toxicity of the compound, since CCS did not affect splenocyte or GA15 cell viability. The viability of the cells was not significantly affected by culture in the presence of CCS for periods of time greater than 48 hours.

25 Since we have shown that fraction CCS from crude bromelain blocks activation of the MAP kinase pathway and blocks T cell activation, CCS may be of use in the treatment of T cell-mediated diseases.

In addition to its importance for IL-2 production and T cell activation, the MAP kinase pathway is also important for the production of growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF) and insulin-like growth factor (IGF). CCS will therefore block the production of these, and
5 other, growth factors and the production of other cytokines such as IL-4, IFN- γ , GM-CSF and many more.

Also, as briefly mentioned above, CCS is likely to be of use in the treatment of
10 cancer.

Thus, CCS may be of use in a method for the treatment or prevention of diseases or conditions mediated by:

- i. activation of T cells;
 - 15 ii. activation of the MAP kinase pathway; or
 - iii. the production of growth factors or cytokines;
- or in the treatment or prevention of cancer.

In a second aspect of the invention, therefore, there is provided a component of
20 bromelain which contains proteins having molecular weights of about 15.07kDa, 25.85kDa and 27.45kDa as determined by SDS-PAGE, has isoelectric points of 10.4 and 10.45 and is obtainable by the following method:

- i. dissolving bromelain in acetate buffer at pH 5.0;

- 25 ii. separating the components of the bromelain by fast flow high performance liquid chromatography on S-sepharose eluting with a linear gradient of 0 to 0.8 M sodium chloride in acetate buffer over 300 ml;

iii. collecting the fraction corresponding to the final double peak off the column; and

iv. isolating the protein from the fraction collected in (iii)

5

for use in medicine, particularly in the treatment or prevention of diseases and conditions mediated by:

- i. activation of T cells;
 - ii. activation of the MAP kinase pathway; or
 - 10 iii. the production of growth factors or cytokines;
- or in the treatment or prevention of cancer.

On further analysis of the CCS fraction of bromelain, the present inventors have found that it comprises more than one component. Sequencing of the proteins in the fraction
15 showed that it consists of the cysteine proteases ananain and comosain together with various other components.

Thus, it appears that both ananain and comosain or a mixture of the two may be responsible for the activity of the CCS fraction of bromelain.

20

In a further aspect of the invention, therefore, there is provided the use of ananain, comosain, a mixture of ananain and comosain or the CCS fraction of bromelain in the preparation of an agent for treatment or prevention of diseases or conditions mediated by:

- 25 i. activation of T cells;
 - ii. activation of the MAP kinase pathway; or
 - iii. the production of growth factors or cytokines;
- or in the treatment or prevention of cancer.

In our earlier application WO-A-9600082 we discussed the inhibition of the MAP kinase cascade by crude bromelain. However, at that time, we were not able to determine which component of the crude bromelain mixture was responsible for this activity although we speculated that it might be stem bromelain protease. We have now discovered that in addition to blocking cyclic nucleotide pathways, stem bromelain protease does have some activity against the MAP kinase pathway. However, it is far less effective in blocking the MAP kinase cascade than the CCS fraction of bromelain of the present invention. Indeed, we have now found that the CCS fraction of bromelain is in the region of ten orders of magnitude more active than stem bromelain protease in blocking MAP kinase activation.

The activation of the MAP kinase pathway in T cells to produce IL-2 and drive T cell clonal expansion is an essential component of the immune response. The absence of this process can have fatal consequences, as can be observed in people with AIDS or genetic mutations which result in T cell defects. However, the activation of T cells can also lead to detrimental consequences. For example, if autoreactive T cells are activated, autoimmune diseases can result. CCS is therefore likely to be of use in the treatment of autoimmune diseases such as rheumatoid arthritis, type-1 diabetes mellitus, multiple sclerosis, Crohn's disease and lupus.

Also, the activation of T cells specific for engrafted tissue can lead to graft or transplant rejection and so CCS may also be of use in preventing this.

The activation of allergen-specific T cells can cause allergic reactions. Inflammatory cytokines and other cellular products, such as histamine, are released from cells following exposure to allergens. The release of histamine and inflammatory cytokines involves the MAP kinase pathway and so blocking of the MAP kinase

pathway with CCS is likely to be an effective treatment for allergies.

In addition, CCS is likely to be of use in the prevention of toxic shock and other diseases mediated by over production of bacterial endotoxins. Toxic shock is mediated by the production of lipopolysaccharides (LPS) from gram-negative bacteria. LPS triggers the production of TNF- α and interleukin-1 *via* activation of the MAP kinase pathway in macrophages. The secretion of these cytokines elicits a cascade of cytokine production from other cells of the immune system (including T cells), which leads to leucocytosis, shock, intravascular coagulation and death.

A further use for CCS is in the prevention of programmed cell death (apoptosis). This is a special event whereby cells are stimulated to destroy their own DNA and die. It is an essential event in most immune responses (to prevent the accumulation of too many cells), but can also have immunosuppressive consequences in some instances, such as in HIV infection and ageing so that too many cells die and there are insufficient left to combat infection (Perandones *et al*, 1993, *J. Immunol.*, **151**, 3521-3529). Because the initiation of apoptosis is dependent on specific cell signalling events, including activation of the MAP kinase pathway, CCS is likely to be effective in blocking apoptosis.

The continual activation of T cells during chronic disease can also lead to pathological consequences, as can be found in certain chronic parasitic infections, such as chronic granulomas diseases such as tuberculoid leprosy, schistosomiasis and visceral leishmaniasis. Furthermore, the invasion of parasites and pathogens, and their subsequent survival in cells, is dependent on these organisms utilising host cell signalling pathways (Bliska *et al*, 1993, *Cell*, **73**, 903-920). For example, *Salmonella* has been demonstrated to phosphorylate MAP kinase, which allows for the bacteria to become endocytosed by macrophages (Galan *et al*, 1992, *Nature*,

357, 588-589). The bacteria then proliferate and destroy the cell. Because CCS has been shown to modify host signalling pathways, and, in particular, to inhibit MAP kinase, another of its potential applications could be to inhibit invasion by parasites and pathogens and their survival in cells.

5

CCS may also be of use for the treatment of cancer and, indeed, we have shown that CCS can block human tumour growth *in vitro*. The anti-tumour mechanism of action of CCS remains to be determined but seems likely to be a result of the blocking of activation of the ERK-2 pathway.

10

As mentioned earlier, MAP kinase activation is dependent on p21^{Ras} and Raf-1, which are important oncogenes. p21^{Ras} and Raf-1 proteins help to relay signals from growth factor receptors on the surface of cells to MAP kinases to stimulate cell proliferation or differentiation. Oncogenic (or mutant) p21^{Ras} or Raf-1 genes produce defective proteins that have acquired independence from externally supplied growth factors and, at the same time, may no longer respond to external growth-inhibitory signals. Mutant p21^{Ras} or Raf-1 proteins are thus persistently hyperactive and their unbridled catalytic activity has a detrimental effect on the control of cell growth. Oncogenic p21^{Ras} or Raf-1 genes therefore promote cancer and tumour formation by disrupting the normal controls on cell proliferation and differentiation. Approximately 30% of human cancers have mutations in a p21^{Ras} gene.

15

20

25

Given that signals transmitted by p21^{Ras} and Raf-1 can be blocked *via* MAP kinase, CCS would be expected to block cancer and tumour growth. The protein fraction of the present invention would therefore be useful for treating many different types of cancer including solid cancers such as ovarian, colon, breast or lung cancer and melanoma as well as non-solid tumours and leukaemia.

The bromelain fraction of the invention will usually be formulated before administration to patients and so, in a further aspect of the invention there is provided a pharmaceutical or veterinary composition comprising the CCS fraction of bromelain together with a pharmaceutically or veterinarily acceptable excipient.

5

The CCS fraction may be administered by a variety of routes including enteral, for example oral, nasal, buccal, topical or anal administration or parenteral administration, for example by the intravenous, subcutaneous, intramuscular or intraperitoneal routes.

10

In many cases, the oral route may be preferred as this is often the route which patients find most acceptable. The oral route may be particularly useful if many doses of the protein are required.

15

When oral administration is chosen, it may be desirable to formulate the CCS fraction in an enteric-coated preparation in order to assist its survival through the stomach. Alternatively, another orally administrable dosage form may be used, for example a syrup, elixir or a hard or soft gelatin capsule, either of which may be enteric coated.

20

However, under certain circumstances, it may more convenient to use a parenteral route. For parenteral administration, the protein may be formulated in distilled water or another pharmaceutically acceptable solvent or suspending agent.

25

A suitable dose of the CCS fraction to be administered to a patient may be determined by the clinician. However, as a guide, a suitable dose may be from about 0.5 to 20 mg per kg of body weight. It is expected that in most cases, the dose will be from about 1 to 15 mg per kg of body weight and preferably from 1 to 10 mg per kg of body weight. For a man having a weight of about 70 kg, a typical dose would therefore be from about 70 to 700 mg.

The invention will now be further described with reference to the following examples and to the drawings in which:

- 5 FIGURE 1 is a diagrammatic representation of signal transduction events associated with T cell activation that lead to IL-2 production.

FIGURE 2 is an ultra violet elution profile of crude bromelain after cation exchange chromatography on SP Sepharose high performance media.

10

FIGURE 3 is a plot showing the proteolytic activity and the protein content of crude bromelain fractions after cation exchange chromatography on SP Sepharose high performance media.

- 15 FIGURE 4 is an SDS-PAGE of SP Sepharose high performance chromatography pooled fractions run on 4-20% T gradient gels with lanes 1 to 4 and 6 to 9 containing proteins CCT, CCV, CCX and CCZ and CCY, CCW, CCU and CCS respectively and lanes 5 and 10 containing molecular weight markers.

- 20 FIGURE 5 shows isoelectric focussing of pooled fractions run on pH 3-11 gradient gels with Lanes 1, 11 and 12 showing high IEF markers, Lanes 2 and 13 showing crude bromelain and Lanes 3 to 10 showing proteins CCT, CCV, CCX, CCZ, CCY, CCW, CCU and CCS respectively.

- 25 FIGURE 6 is a Western blot using anti-phosphotyrosine mAb which demonstrates that CCS reduces tyrosine phosphorylation of p42 kDa (ERK-2) protein. Th0 cells were treated with bromelain fractions (50µg/ml) for 30 min, washed and then stimulated with combined PMA (20 ng/ml) and ionophore (1µM) for 5 min. Unstimulated cells

served as controls. Cells were then lysed and postnuclear supernatants were subjected to SDS-PAGE and Western blotting. In this figure, closed symbols indicate proteins phosphorylated by combined PMA plus ionophore. Open symbols indicate ERK-2 protein reduced by CCS treatment.

5

FIGURE 7 is a Western blot using anti-phosphotyrosine mAb which demonstrates that CCS increases tyrosine phosphorylation of protein substrates. Th0 cells were treated with CCS, crude bromelain (Brom), stem bromelain protease (SBP) or CCT fraction (50µg/ml) for 30 min, washed and then stimulated with combined PMA (20 ng/ml) and ionophore (1µM) for 5 min. Unstimulated cells served as controls (Cont). Cells were then lysed and postnuclear supernatants were subjected to SDS-PAGE and Western blotting. In this figure, closed symbols indicate proteins phosphorylated by CCS but not by other treatments. Open symbols indicate phosphoproteins protein reduced by CCS and crude bromelain treatment.

15

FIGURE 8 is a Western blot using anti-phosphotyrosine mAb which shows that the inhibitory effect of CCS on ERK- 2 is dependent on its proteolytic activity and occurs in a dose-dependent manner. Th0 cells were treated with CCS (0 to 25µg/ml) or CCS incubated with the selected protease inhibitor, E-64, for 30 min. Cells were then washed and then stimulated with combined PMA (20 ng/ml) and ionophore (1µM) for 5 min. Cells were then lysed and postnuclear supernatants were subjected to SDS-PAGE and Western blotting. In this figure, closed symbols indicate proteins phosphorylated by CCS. Open symbols indicate ERK-2 phosphoprotein inhibited by active CCS but not by inactivated CCS.

20

FIGURE 9 is an immunoblot which confirms that the 42kDa phosphoprotein inhibited by CCS is ERK-2. Th0 cells were treated with CCS (50µg/ml) for 30 min, or untreated, washed and then stimulated with combined PMA (20 ng/ml) and ionophore

25

(1 μ M) for 5 min. Cell lysates were immunoblotted with anti-ERK-2 mAb

FIGURE 10 is a Western blot using anti-phosphotyrosine mAb which shows that crosslinked anti-CD3 ϵ mAb induces tyrosine phosphorylation of multiple proteins. Th0
5 cells were stimulated with crosslinked anti-CD3 ϵ for 0 to 20 min. Cells were then lysed and postnuclear supernatants were subjected to SDS-PAGE and Western blotting. Closed symbols denote anti-CD3 ϵ mAb-induced tyrosine phosphorylated proteins.

10 FIGURE 11 is a Western blot using anti-phosphotyrosine mAb which demonstrates that CCS inhibits tyrosine phosphorylation in TCR-stimulated T cells. Th0 cells were treated with CCS (0 to 5 μ g/ml) for 30 min, washed and then crosslinked anti-CD3 ϵ mAb for 5 min. Cells were then lysed and postnuclear supernatants were subjected to SDS-PAGE and Western blotting. In this figure, the symbols denote anti-CD3 ϵ mAb-
15 induced tyrosine phosphorylation of ERK-2, which is reduced by CCS.

FIGURE 12 is a Western blot using anti-Raf-1 mAb which shows that CCS inhibits the mobility shift of Raf-1. . Th0 cells were treated with CCS (0 to 50 μ g/ml) for 30 min, washed and then stimulated with either (A) combined PMA plus ionophore or (B) then
20 crosslinked anti-CD3 ϵ mAb for 5 min. Cells were then lysed and postnuclear supernatants were subjected to SDS-PAGE and Western blotting.

FIGURE 13 is a pair of plots showing that CCS decreases IL-2 production and proliferation in purified CD4⁺ T cells. T cells were treated with CCS (50 μ g/ml),
25 washed and then cultured in either media alone or with immobilised anti-CD3 ϵ mAb and soluble anti-CD28 mAb. (A) IL-2 production was determined by the CTL-L assay as described in Example 5. (B) Proliferation was determined by the incorporation of

^3H -thymidine. CD4^+ T cells cultured in the absence of mAb (stimuli) did not produce any detectable IL-2 and do not proliferate.

FIGURE 14 is a pair of plots showing that CCS decreases IL-2 production by splenocytes but does not inhibit splenocyte proliferation. Splenocytes were treated with CCS (50 $\mu\text{g/ml}$), washed and then cultured in either media alone or with immobilised anti-CD3 ϵ mAb. (A) IL-2 production was determined by the CTL-L assay as described in Example 5. (B) Proliferation was determined by the incorporation of ^3H -thymidine. Splenocytes cultured in the absence of mAb (stimuli) did not produce any detectable IL-2 and do not proliferate).

FIGURE 15 is a plot which shows that CCS inhibits tumour cell growth *in vitro*. Cancer cell lines were treated with CCS (50, 10, 2.5, 1 and 0.25 $\mu\text{g/ml}$) or water as a control. After 96 h treatment, the effect of CCS on tumour cell growth was evaluated. Columns represent the 50% inhibitory concentrations (IC_{50} $\mu\text{g/ml}$) of CCS (the amount of CCS required to inhibit 50% of tumour cell growth).

EXAMPLE 1 - Purification of Bromelain Proteins

a. Materials

Reagents Bromelain (E.C 3.4.22.4; proteolytic activity, 1,541 nmol/min/mg) was obtained from Solvay Inc. (Germany). Fast Flow S Sepharose, Pharmalyte 3-10TM, Ampholine 9-11TM, Ready Mix IEFTM (acrylamide, bisacrylamide) and IEFTM markers were obtained from Pharmacia Biotech. Precast 4-20% acrylamide gels and broad range molecular weight markers were obtained from Bio-Rad Laboratories. All other reagents were of analytical grade and obtained from either Sigma Chemical Co. or British Drug House.

b. Proteinase Assay

The proteolytic activity of bromelain was determined by use of an in-house microtitre

plate based assay using the synthetic substrate Z-Arg-Arg-pNA. This assay was based on that described by Filippova *et al* in *Anal. Biochem.*, **143**, 293-297 (1984). The substrate was Z-Arg-Arg-pNA as described by Napper *et al* in *Biochem. J.*, **301**, 727-735, (1994).

5

c. **Protein Assay**

Protein was measured using a kit supplied by Bio-Rad that is a modified method of Lowry *et al* (*J. Biol. Chem.* (1951) **193**, 265-275). Samples were compared to bovine serum albumin standards (0 to 1.5 mg/ml) prepared in either 0.9 % saline or 20 mM acetate buffer pH 5.0, as appropriate.

10

d. **Preparation of Bromelain**

All the following steps were performed at ambient temperature (20 to 25°C). A solution of bromelain (30 mg/ml) was prepared by dissolving 450 mg of powder in 15 ml of 20 mM acetate buffer (pH 5.0) containing 0.1 mM EDTA, sodium. The solution was dispensed into 10 x 1.5 ml microcentrifuge tubes and centrifuged at 13,000 x g for 10 minutes to remove insoluble material. The clear supernatants were pooled and used for chromatography

15

e. **Fast Flow S-Sepharose High Performance Chromatography**

A Fast flow S-sepharose column was prepared by packing 25 ml of media into an XK 16/20™ column (Pharmacia Biotech) and equilibrated with 20 mM acetate buffer (pH 5.0) containing 0.1 mM EDTA on an FPLC™ system at 3ml/min. 5 ml of bromelain solution was injected onto the column. Unbound protein was collected and the column washed with 100 ml of acetate buffer. Protein bound to the column was eluted with a linear gradient of 0 to 0.8 M NaCl in acetate buffer over 300 ml. 5 ml fractions were collected throughout the gradient and Figure 2 shows a typical U.V. chromatogram of crude bromelain obtained from this procedure.

20
25

5 The fractions were then analysed for protein and proteolytic activity as described above and Figure 3 shows the proteolytic activity against the synthetic peptide Z-Arg-Arg-pNA and the protein content of the individual fractions. The protein content profile closely mirrors that of the U.V., as expected, but the main proteolytic activity is confined to the two major peaks that correspond to that of bromelain protease (SBP). Small activities are observed in other areas of the chromatogram that may corresponds to other proteases distinct from SBP, such as the later eluting CCS fraction, which contains ananain and comosain.

10

The main peaks identified from the U.V. profile were pooled from three successive runs and named as indicated in Table 1. Pooled fractions were used for physico-chemical characterisation. Pooled fractions were concentrated by ultrafiltration and buffer exchanged using PD10 columns into isotonic saline (0.9 % w/v NaCl). The protein content and Z-Arg-Arg-pNA activity were calculated prior to biological testing and are shown in Table 2.

15

The pooled fractions were processed for analysis as described below.

20

Table 1. Summary of Pooled Fractions from SP Sepharose HP Fractionated Bromelain (QC2322)

Component	Description	Fractions Pooled (Inclusive)
CCT	Flow through (unbound components)	Unbound column flow through
CCV	First peak off column	8-9
CCX	Second sharp peak off column	13-14
CCZ	Small peak on ascending edge of the third main bromelain peak	19-20
CCY	First main bromelain peak	23-24
CCW	Second main bromelain peak	27-29
CCU	Small peak on descending edge of the second main bromelain peak	33-34
CCS	Last double peak off column	39-44

5 **Table 2. Calculated Protein Content and Z-Arg-Arg-pNA Activity of Pooled Fractions used for Testing Biological Activity.**

Pooled Fractions	Z-Arg-Arg-pNA Activity (μ Moles/min/ml)	Protein Content (mg/ml)
CCT	11.30	1.00
CCV	9.78	1.00
CCX	71.71	1.00
CCZ	688.81	1.00
CCY	1500.0	0.574
CCW	1500.0	0.543
CCU	1500.0	0.421
CCS	379.76	1.00

f. **Processing of Pooled Fractions**

The proteolytic activity and protein content of pooled fractions were determined and the concentrations adjusted to approximately either 1.4 mg/ml of protein or 105
5 nmoles/min/ml of proteinase activity using a Filtron™ stirred cell containing an ultrafiltration membrane of nominal molecular weight cut-off of 10 kDa. The fractions were then buffer exchanged using PD10™ columns (Pharmacia Biotech) into isotonic saline (0.9 % w/v NaCl), sterile filtered (0.2 µm) and adjusted for protein content or
10 proteolytic activity. Samples were then frozen at -80°C and used in the *in vitro* studies described below.

g. **Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis (SDS-PAGE)**

Pooled FPLC™ samples were analysed by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) on precast 4 to 20 % T gradient gels. Samples were
15 prepared for electrophoresis by acid precipitation in which 100 µl was mixed with an equal volume of 20 % w/v trichloroacetic acid (TCA). Precipitated protein was collected by centrifugation at 13,000 x g for 10 minutes and the supernatant discarded. The pellet was washed twice with 0.5 ml of diethyl ether and left to dry in air at ambient temperature. The pellets were then dissolved in 300 µl of SDS-PAGE sample
20 buffer (62.5 mM Tris-HCl pH 6.8 containing 10 % v/v glycerol, 2 % w/v sodium dodecyl sulphate and 40 mM dithiothreitol) and heated at 95°C in a water bath.

SDS-PAGE broad range molecular weight standards diluted 1:20 in SDS-PAGE sample buffer were treated similarly and run with the samples. Gels were run on a mini
25 Protean II™ electrophoresis system according to Bio-Rad's protocol at 240 V and until the dye front reached the end of the gel (30 to 45 min).

After electrophoresis, separated proteins were stained overnight with orbital mixing in

a solution of 0.075 % w/v colloidal brilliant blue G-250 containing 1.5 % v/v phosphoric acid, 11.25% w/v ammonium sulphate and 25 % v/v methanol. Gels were destained, to obtain a clear background, in a solution of 25 % v/v methanol and 10 % v/v acetic acid.

5

RESULTS

The purity of fractions is shown by SDS-PAGE in Figure 4. All of the pooled fractions except the column flow through (CCT) showed that the major protein present was of molecular weight between approximately 25-28 kDa. This corresponds to the molecular weight of cysteine proteinases isolated from bromelain by other authors (Rowan *et al*, *Methods in Enzymology*, (1994), **244**, 555-568). The purity of fractions CCX, CCZ, CCY and CCW appears to be high. Minor components of lower molecular weight can be observed in some fractions, particularly CCT, CCV, CCX and CCS. Pooled fractions CCU and CCS contain a doublet between 25-28 kDa; the higher gel loadings of fractions CCX, CCZ, CCY, and CCW means that doublet bands may also be present in these fractions. A summary of the components and their calculated molecular weights in pooled fractions, as determined by SDS-PAGE, is shown in Table 3.

Proteins in pooled fractions CCX, CCZ, CCY+CCW and CCU were transferred onto nitro-cellulose after SDS-PAGE by Western blotting and probed with rabbit antisera raised against purified stem bromelain protease (SBP) (results not shown). All protein bands in these pooled fractions were recognised by antibodies in the sera, indicating immunologically similar proteins, probably belonging to the cysteine proteinase family of enzymes.

Table 3. Summary of the Molecular Weights of Proteins found in SP Sepharose HP Pooled Fractions as Determined by SDS-PAGE.

Pooled Fractions	Molecular Weight (kDa) of Major Protein Band(s)	Molecular Weight (kDa) of Minor Protein Band(s)
CCT	76.03	15.07
CCV	15.07, 25.85, 28.28, 76.03	
CCX	25.08	15.07, 76.03
CCZ	27.45	13.37, 16.49, 76.03
CCY	27.45	6.5
CCW	27.45	
CCU	27.45, 28.28	
CCS	15.07, 25.85, 27.45	

5 h. **Isoelectric Focusing**

Pooled fractions (0.5 to 1.0 mg/ml) were diluted 1:3 with deionised water and run on gradient gels of pH 3 to 11. Gels were cast using Ready Mix IEF™ to produce a 5.5 % T, 3 % C polyacrylamide gel containing 10 % v/v glycerol, 5.0 % Pharmalyte 3-10™ and 2.5 % Ampholine 9-11™. Briefly, 10 µl of sample and high pI markers were loaded onto the gel after prefocusing at 700 V. Sample entry was at 500 V for 10 min, focusing was at 2500 V for 1.5 hour and band sharpening at 3000 V for 10 min. After electrophoresis the proteins were fixed with a solution of 20 % w/v TCA for 30 min, washed in destain for 30 min to remove TCA and stained with brilliant blue G-250 as described for SDS-PAGE (see above).

15

RESULTS

Figure 5 shows that all fractions except CCX contained basic proteins focusing beyond the 9.3 pI marker. Localised charge interactions with the chromatographic media functional groups may explain why proteins of pI 3.8 and 3.85 in CCX, adsorbed onto a cation exchange resin at pH 5.0. CCZ was present as a single band of pI 9.7, whilst pooled fractions CCY, CCW, and CCU contained multiple bands of isoelectric points in the range pH 9.5-9.8. At least part of this heterogeneity can be explained by variation in the carbohydrate moiety of a common stem bromelain protein backbone. The values are in agreement with those reported in the literature of pI 9.45-9.55 for bromelain (Rowan *et al*, *Methods in Enzymology*, (1994), **244**, 555-568). Pooled fractions CCS contains two basic proteins of pI greater than 10.25. Estimates by extrapolation give pIs of 10.4 and 10.45. These correspond to ananain and comosain, and are in agreement with other estimates (Rowan *et al*, as above) of pIs greater than 10. The pIs of proteins in each of the pooled fractions are summarised in Table 4.

Table 4. Summary of the estimated Isoelectric points of Proteins found in SP Sepharose HP Pooled Fractions.

Pooled Fractions	Isoelectric Points of Proteins
CCT	Not detected
CCV	Not Detected
CCX	3.8, 3.85
CCZ	9.7
CCY	9.6, 9.7
CCW	9.57, 9.6, 9.7
CCU	9.57, 9.6, 9.75
CCS	10.4, 10.45

EXAMPLE 2 - NH₂-terminal Amino Acid Analysis of Bromelain Components

In a separate experiment, pooled fractions of bromelain were run by SDS PAGE and blotted as above onto PVDF membrane. The membrane was stained with 0.025 % w/v coomassie blue R-250, dissolved in 40 % v/v methanol for 10 min, followed by destaining in 50 % v/v methanol. The membrane was dried in air at room temperature and NH₂-terminal amino acid sequencing of the stained proteins was carried out. Briefly, the protein band was cut from the membrane and placed in the upper cartridge of the sequencer. NH₂-terminal amino acid analysis of bromelain components was determined by Edman degradation using a gas phase sequencer (Applied Biosystems), equipped with an on-line phenylthiohydantion amino acid analyser. Table 5 shows the first 21 NH₂-terminal amino acids of CCZ, CCZ, stem bromelain protease, ananain and comosain.

Table 5. NH₂-Terminal Sequence Similarities of CCZ Protein and Those of Known Proteinases Isolated from Bromelain.

Proteinase	Position from N-Terminus
CCZ	V L P D S I D W R Q K G A V T E V K N R G 1 5 10 15 20
CCX	V P Q S I D W R D Y G A V N E V K N 4 9 14
Stem Bromelain Protease	A V P Q S I D W R D Y G A V T S V K N Q N 1 5 10 15 20
Ananain	V P Q S I D W R D S G A V T S V K N Q G 1 4 9 14 19
Comosain	V P Q S I D W R N Y G A V T S V K N Q G 1 4 9 14 19

All proteins share sequence homologies. Ananain and comosain differ by 2 out of 20 amino acids when compared to stem bromelain protease. CCZ differs by 8 out of 21 amino acids when compared to stem bromelain protease. CCZ differs from ananain and comosain by 6 out of 20 amino acids. Comosain differs by 2 amino acids from ananain. Whilst it is clear that these proteins are structurally related, they are all distinct, showing divergence from each other. These proteinases also differ in their proteinase substrate specificity and their biological activity.

EXAMPLE 3 - Fraction CCS inhibits tyrosine phosphorylation of p42 kda phosphoprotein.

a. Materials

Antibodies Anti-CD3 ϵ -chain mAb (145-2C11) and anti-CD28 mAb (PV-1) were purchased from Pharmingen (San Diego, CA) and goat anti-hamster IgG Ab was from Sigma (Dorset, UK). Mouse anti-phosphotyrosine mAb (4G10), mouse anti-MAPk R2 (ERK-2) mAb and mouse anti-Raf-1 mAb were from UBI (Lake Placid, NY). Goat anti-mouse and goat anti-rabbit IgG Ab conjugated to horse radish peroxidase (HRP) were from BioRad (Hemel Hemstead, Hertfordshire, UK). Rabbit polyclonal phospho-specific MAPk IgG which recognise tyrosine phosphorylated p44 and p42 MAPks were from New England BioLabs (Hitchin, Hertfordshire, UK).

Reagents Phorbol 12-myristate 13-acetate (PMA) and calcium ionophore A23187 were purchased from Sigma. Bromelain (E.C 3.4.22.4; proteolytic activity, 1,541 nmol/min/mg) was obtained from Solvay Inc. (Germany). E-64 (L-trans-epoxysuccinylleucylamido-(4-guanidino)butane, a selective cysteine protease inhibitor, was from Sigma.

Cells The T cell hybridoma GA15 was a generous gift from B. Fox (ImmuLogic Pharmaceutical Corporation, Boston, MA). GA15 was generated by fusing the

thymoma BW5147 with the T_H2 clone F4 specific for KLH in association with I-A^b, and were maintained as previously described (Fox, 1993, *Int. Immunol.*, 5, 323-330). GA15 exhibit a T_H0 cell phenotype as they produce IL-2, IL-4 and IFN- γ following stimulation with crosslinked anti-CD3 ϵ mAb (Fox, 1993).

5

b. **Stimulation of T cells.** Cells (2×10^7) suspended in RPMI 1640 were treated with CCS (1 to 50 $\mu\text{g/ml}$) diluted in saline (0.9% (w/v)) for 30 min at 37°C. Mock treated cells were treated with an equal volume of saline (diluent). At high concentrations of CCS (50 or 100 $\mu\text{g/ml}$) cell aggregation occurred, as noted previously in studies with crude bromelain. Following treatment, cell aggregates were gently dispersed by washing cells 3 times and then resuspending in fresh RPMI. Cells were stimulated via the cell surface with crosslinked mAb to the TCR (anti-CD3 ϵ), or directly, using combined PMA (20 ng/ml) and ionophore (1 μM) for times indicated in figure legends and the text.

15

Stimulation via the TCR was conducted by first incubating T cells on ice for 30 min with anti-CD3 ϵ mAb (20 $\mu\text{g/ml}$). Excess mAb was then removed by washing once at 4°C and anti-CD3 ϵ mAb was crosslinked with goat anti-hamster IgG (20 $\mu\text{g/ml}$) at 37°C. Stimulation was terminated by the addition of ice-cold lysis buffer (25 mM Tris, pH 7.4, 75 mM NaCl, 2 mM EDTA, 0.5% Triton X-100, 2 mM sodium orthovanadate, 10 mM sodium fluoride, 10 mM sodium pyrophosphate, 74 $\mu\text{g/ml}$ leupeptin, 740 μM PMSF and 74 $\mu\text{g/ml}$ aprotinin) for 30 min with continual rotation at 4°C. Lysates were clarified (14,000 x g for 10 min) and an equal volume of 2 X SDS-PAGE sample buffer (50 mM Tris, pH 7, 700 mM 2-ME, 50% (v/v) glycerol, 2% (w/v) SDS, 0.01% (w/v) bromophenol blue) was added to postnuclear supernatants. Proteins were solubilised at 100°C for 5 min and samples containing 1×10^6 cell equivalents were resolved by SDS-PAGE.

20
25

c. **Immunoblotting.** Separated proteins were transferred to nitrocellulose membranes (Bio-Rad) which were then blocked with 5% (w/v) bovine serum albumin (Sigma, fraction V; BSA), 0.1% Nonidet P-40TM in Tris-buffered saline (170 mM NaCl and 50mM Tris, pH 7.4; TBS). Immunoblots were incubated with the appropriate antibodies as indicated in figure legends. Primary antibodies were diluted in antibody dilution buffer comprised of 0.5% (w/v) BSA, 0.1% (v/v) Tween-20 in TBS at 4 °C for 2 h followed by detection with the appropriate secondary antibody conjugated to horseradish peroxidase diluted in antibody dilution buffer at 4 °C for 1 h. Following each incubation step, membranes were washed extensively with 0.1% Tween-20 in TBS. Immunoreactivity was determined using the ECL chemiluminescence detection system (Amersham Corp., Arlington Heights, IL).

d. **Inhibition of proteolytic activity of CCS.** A specific cysteine protease inhibitor, E-64, was used to inactivate the proteolytic activity of CCS. CCS (25 µg/ml) diluted in 3 µM dithiothreitol, 100 µM E-64, 60 mM sodium acetate (pH 5) was incubated for 10 minutes at 30°C. The inactivated CCS was then dialysed overnight in saline at 4°C. Earlier studies with crude bromelain have shown that these conditions are sufficient to induce 99.5% inactivation of proteolytic activity as assayed with the Z-Arg-Arg-pNA substrate (see above). T cells were treated with E-64 inactivated CCS (25 µg/ml) and compared with untreated CCS and mock-treated T cells stimulated with PMA plus ionophore.

RESULTS

a. **Fraction CCS inhibits tyrosine phosphorylation of p42 kda phosphoprotein.** We have previously shown that bromelain blocks tyrosine phosphorylation of ERK-2 following stimulation of T cells with combined PMA plus calcium ionophore (WO-A-96/00082). Phorbol ester and ionophore stimulation of T

cells act synergistically to reproduce many features of TCR stimulation such as IL-2 secretion, IL-2 receptor expression, and T cell proliferation (Truneh *et al.*, 1985, *Nature*, **313**, 318-320; Rayter *et al.*, 1992, *EMBO*, **11**, 4549-4556). Phorbol esters can mimic antigen receptor triggering and bypass TCR-induced protein tyrosine
5 kinases to activate ERK-2 by a direct agonist action on PKC and p21^{Ras}. Calcium ionophore A23187 induces increased intracellular release of Ca²⁺ and therefore mimics the action of inositol 1,4,5-trisphosphate (IP₃). Phorbol esters and ionophore however, stimulate PKC pathways that are not controlled by the TCR (Izquierdo *et al.*, 1992, *Mol. Cell. Biol.*, **12**, 3305-3312) suggesting separate intracellular
10 pathways within T cells that regulate T cell function. We therefore investigated which fraction of bromelain could block T cell signalling via the TCR-independent pathway by examining its effect on PMA and ionophore-induced tyrosine phosphorylation.

15 Stimulation of T cells with combined ionophore and PMA induced tyrosine phosphorylation of several proteins including those of circa 100 kda, 85 kda, 42 kda and 38 kda. CCS (50 µg/ml) pre-treatment reduced tyrosine phosphorylation of the p42 kda protein, and did not significantly inhibit phosphorylation of any other substrate (Fig 6). In two experiments, CCS, but no other fraction, also increased
20 tyrosine phosphorylation of circa 36 kda, 38 kda, 85 kda, 94 kda and 102 kda proteins (Fig 7 and Fig 8).

The ability of CCS to block tyrosine phosphorylation of the 42 kda phosphoprotein was dose-dependent (Fig 8) and dependent on its proteolytic activity, since E-64
25 completely abrogated the inhibitory effect of CCS on p42 kda phosphorylation (Fig 8). E-64 treatment of T cells did not affect PMA and ionophore-induced T cell signalling.

CCS inhibits ERK-2 tyrosine phosphorylation. We suspected that the 42 kda phosphoprotein inhibited by CCS was the MAPk ERK-2, so we conducted immunoblot analysis with specific anti-ERK-2 mAb and anti-phospho MAPk antibodies, which specifically detects ERK-1 and ERK-2 only when catalytically
5 activated by phosphorylation at Tyr204. Immunoblotting of CCS-treated cells that had been stimulated with PMA plus ionophore, confirmed that the p42 kda phosphoprotein was indeed ERK-2 (Fig 9).

CCS reduces TCR-induced tyrosine phosphorylation of ERK. We next
10 investigated the effect of CCS on TCR-mediated signal transduction by assessing substrate tyrosine phosphorylation of GA15 stimulated with crosslinked anti-CD3ε mAb. Immunoblots of GA15 lysates, using specific anti-phosphotyrosine mAb, revealed increased tyrosine phosphorylation of multiple proteins including those of circa 120 kda, 100 kda, 85 kda, 76 kda, 70 kda, 42 kda and 40 kda, consistent with
15 phosphoproteins observed in other T cell lines following TCR-ligation (June *et al*, 1990, *J. Immunol.*, **144**, 1591-1599 and *Proc. Natl. Acad. Sci. USA*, **87**, 7722-7726, reviewed by Cantrell, 1996, *Annu. Rev. Immunol.*, **14**, 259-274) (Fig 10). Tyrosine phosphorylated proteins were readily detected between 2 and 5 min following stimulation and remained phosphorylated for at least 10 min (Fig 10).
20 GA15 cells stimulated with anti-CD3 mAb alone or cross-linking Ab, did not induce tyrosine phosphorylation of any cellular substrate (data not shown). Again, CCS pre-treatment of GA15 for 30 min caused a reduction in TCR-induced protein tyrosine phosphorylation of ERK-2 in a dose-dependent manner (Fig 11). CCS did not markedly affect tyrosine phosphorylation of other TCR-induced phosphoproteins,
25 suggesting that CCS has a selective mode of action.

EXAMPLE 4 – CCS retards the mobility shift of Raf-1

Raf-1 is an immediate upstream activator of MEK-1 which activates ERK-2. Raf-1

activation requires phosphorylation on specific serine and threonine residues (Avruch et al., 1994, *TIBS*, 19, 279-283). To investigate whether CCS affects any other substrates upstream from ERK-2 in the MAP kinase cascade, we investigated the effect of CCS on Raf-1. T cells were treated with CCS (0 to 50 µg/ml) and then
5 stimulated with either anti-CD3ε mAb or combined PMA plus ionophore as described earlier. Results show that CCS blocks the mobility shift of Raf-1, indicating that it blocks its protein phosphorylation and thus activation. This data confirms that CCS has an effect on the MAP kinase cascade (Fig 12) and that the effect of CCS may not be directly on ERK-2, but on upstream substrates in the
10 MAPk cascade.

EXAMPLE 5 – Effect of CCS on IL-2 Production and T cell Proliferation

a. Materials

Cells Splenocytes were isolated from female BALB/c mice (6-8 weeks old), as
15 previously described in WO-A-96/00082. Highly purified CD4⁺ T cells were isolated from splenocytes using magnetic activated cell sorting (MACS).

b. Interleukin 2 production. T cells diluted in RPMI were treated with CCS (50 µg/ml) or saline at 37°C for 30 min, washed in fresh RPMI and then
20 resuspended in culture medium. T cells were stimulated to produce cytokine mRNA by immobilised anti-CD3ε (4 µg/ml) and soluble anti-CD28 (10 µg/ml). Anti-CD3ε mAb diluted in PBS was immobilised to 24-well, flat bottom, microculture plates (Corning, Corning, NY) by incubation for 16 hours at 4°C. Wells were then washed
three times in PBS prior to addition of triplicate cultures of either splenocytes or
25 purified CD4⁺ T cells (2.5 - 5 x 10⁶ cells per well) which were incubated at 37°C in humidified 5% CO₂ for 24 h. IL-2 levels in the culture supernatant were measured using the CTL-L bioassay (Gillis et al., 1978, *J. Immunol.*, 120, 2027-2032).

c. **T Cell Proliferation.** T cells were treated with CCS (50 µg/ml) for 30 min, washed in RPMI then stimulated with immobilised anti-CD3ε mAb alone or combined anti-CD3ε mAb plus anti-CD28 mAb. Cells were then cultured in 96 well, flat-bottom plates (Nunc) at 10⁵ cells per well for 36 h. Cultures were pulsed with 0.5 µCi of [³H]TdR 12 h prior to harvesting onto glass fibre filters.

RESULTS

a. CCS inhibits IL-2 production and proliferation of CD4⁺ T cells.

Activation of p21^{Ras}, Raf-1, MEK-1 and ERKs are essential for induction of IL-2 transcription in T cells (Izquierdo *et al.*, 1993, *J. Exp. Med.*, **178**, 1199). IL-2 is the major autocrine T cell growth factor which induces proliferation of T cells. The defect in ERK activation demonstrated here could therefore be expected to inhibit IL-2 production and T cell proliferation. We therefore investigated whether CCS could effect a functional outcome of T cell signalling, namely IL-2 production and proliferation in murine splenocytes and highly purified CD4⁺ T cells. CCS (50 µg/ml) treatment of purified CD4⁺ T cells reduced both IL-2 production and proliferation when the ERK pathway was stimulated with anti-CD3ε mAb (Fig 13a and 13b). CCS also blocked IL-2 production by splenocytes, however it did not affect splenocyte proliferation (Fig 14 a and 14b), suggesting that an as yet unidentified component in CCS was acting on accessory cell populations in splenocyte cultures, such as B cells or macrophages. Bromelain can increase costimulatory signals to T cells via an action on B cells. Regardless of the putative effect of CCS on accessory cells, data clearly indicate that CCS blocks IL-2 production and proliferation of purified CD4⁺ T cells, suggesting that CCS blocks T cell activation. IL-2 production and proliferation were dependent on cell stimulation with anti-TCR antibodies as no cytokine was detected in cells cultured in tissue culture media alone (Fig 13 and 14).

EXAMPLE 6 – Effect of CCS on Human Tumour Cell Growth *in vitro***a. Materials**

Cells Tumour cell lines were provided by L. Kelland (Institute of Cancer Research, Sutton, UK) and were as follows; ovarian (SKOV-3, CH-1, A2780), colon (HT29, BE, LOVO), breast (MCF-7, MDA231, MDA361), lung (A549, CORL23, MOR) and melanoma (G361, BOO8, SKMel24).

b. Growth Inhibition of Human Tumour Cell Lines. Studies were conducted by L. Kelland (Institute of Cancer Research, Sutton, UK). Cell lines were trypsinised and single viable cells were seeded into 96-well microtitre plates at a density of 4×10^3 cells/well in 160 μ l growth medium. After allowing for attachment overnight, CCS was then added to quadruplicate wells in 40 μ l of growth medium to give a range of final concentrations in wells of 50, 10, 2.5, 1 and 0.25 μ g/ml. Eight wells served as control, untreated wells. CCS was diluted immediately prior to addition to cells in sterile water. CCS exposure to cells was for 96 h whereupon the cell number in each well was determined by staining with 0.4% sulforhodamine B in 1% acetic acid as described previously (Kelland *et al.*, 1993, *Cancer Res.*, **53**, 2581-2586). 50% inhibitory concentrations (IC₅₀ values in μ g/ml) were then calculated from plots of concentration versus control (%) absorbance (read at 540_{nm}).

RESULTS

a. CCS inhibits human tumour growth *in vitro*. p21^{Ras} and Raf-1 are important oncogenes, which when mutated cause uncontrolled cell growth and proliferation, leading to cancer. Since we have shown that CCS can block the effects of the p21^{Ras}/Raf-1/MEK1/ERK kinase signalling cascade, we investigated whether CCS could block tumour growth. CCS treatment of human tumour cells resulted in a reduction in the growth of several different ovarian, lung, colon, breast and melanoma tumour cell lines *in vitro* (Fig 15). CCS did not affect all cell lines equally, suggesting that CCS has a selective action.

CLAIMS

1. A component of bromelain which contains proteins having molecular weights of about 15.07kDa, 25.85kDa and 27.45kDa as determined by SDS-PAGE, has isoelectric points of 10.4 and 10.45 and is obtainable by the following method:
- 5
- i. dissolving bromelain in acetate buffer at pH 5.0;
 - ii. separating the components of the bromelain by fast flow high performance liquid chromatography on S-sepharose eluting with a linear gradient of 0 to 0.8 M sodium chloride in acetate buffer over 300 ml;
 - 10
 - iii. collecting the fraction corresponding to the final double peak off the column; and
 - 15
 - iv. isolating the protein from the fraction collected in (iii).
2. A component of bromelain as claimed in claim 1, for use in medicine, particularly in the treatment or prevention of diseases and conditions mediated by:
- 20
- i. activation of T cells;
 - ii. activation of the MAP kinase pathway; or
 - iii. the production of growth factors or cytokines;
- or in the treatment or prevention of cancer.
- 25
3. A component of bromelain as claimed in claim 1 for modulating pathways controlling cell growth and proliferation.
4. A component of bromelain as claimed in claim 1 for inhibiting the production

of growth factors or cytokines by cells.

5. A component of bromelain as claimed in claim 1 for inhibiting the activation of the MAP kinase pathway.

5

6. A component of bromelain as claimed in claim 1 for inhibiting the activation of T cells.

10

7. A component of bromelain as claimed in claim 1 for use as an immunosuppressant.

15

8. A component of bromelain as claimed in claim 1 for blocking the production of growth factors and cytokines or for the treatment or prevention of autoimmune diseases, graft or transplant rejection by a host, allergic reactions, toxic shock, apoptosis, parasite or pathogen infections or cancer.

20

9. The use of ananain, comosain, a mixture of ananain and comosain or a component of bromelain as claimed in claim 1 in the preparation of an agent for modulating intracellular signalling pathways which control cell growth and proliferation.

25

10. The use of ananain, comosain, a mixture of ananain and comosain or a component of bromelain as claimed in claim 1 in the preparation of an agent for inhibiting the production of growth factors and cytokines by cells.

11. The use of ananain, comosain, a mixture of ananain and comosain or a component of bromelain as claimed in claim 1 in the preparation of an agent for reducing or preventing the activation of the MAP kinase pathway.

12. The use of ananain, comosain, a mixture of ananain and comosain or a component of bromelain as claimed in claim 1 in the preparation of an agent for reducing or preventing the activation of T cells.

5

13. The use of ananain, comosain, a mixture of ananain and comosain or a component of bromelain as claimed in claim 1, wherein the agent is an immunosuppressive agent.

10 14. The use of ananain, comosain, a mixture of ananain and comosain or the CCS fraction of bromelain in the preparation of an agent for the treatment or prevention of diseases and conditions mediated by

- i. the activation of T cells;
 - ii. activation of the MAP kinase pathway; or
 - 15 ii. the production of growth factors or cytokines;
- or in the treatment or prevention of cancer.

15 15. The use of ananain, comosain, a mixture of ananain and comosain or a component of bromelain as claimed in claim 1 in the preparation of an agent for
20 blocking the production of growth factors and other cytokines or for the treatment or prevention of autoimmune diseases, graft or transplant rejection by a host, allergic reactions, toxic shock, apoptosis, parasite or pathogen infections or cancer.

25 16. A pharmaceutical or veterinary composition comprising the CCS fraction of bromelain together with a pharmaceutically or veterinarily acceptable excipient.


17. A pharmaceutical or veterinary composition as claimed in claim 16, which is adapted for enteral, for example oral, nasal, buccal, topical or anal administration.

18. A pharmaceutical or veterinary composition as claimed in claim 16, which is adapted for parenteral administration, for example by the intravenous, subcutaneous, intramuscular or intraperitoneal routes.

5

19. A pharmaceutical or veterinary composition as claimed in claim 16 which is adapted for oral administration and which is an enteric coated preparation.

1/15



2 / 15




FIG. 2

3 / 15




FIG. 3

4 / 15




FIG. 4

5 / 15





FIG. 5

6 / 15



Ab: α-pY mAb

FIG. 6

7/15


Ab: α -pY mAb

FIG. 7

8 / 15


Ab: α -pY mAb

FIG. 8

9/15




FIG. 9

11 / 15




FIG. 11

12 / 15





FIG. 12

A.

13 / 15



B.





FIG. 13

A.

14 / 15



B.




FIG. 14

15 / 15




FIG. 15

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/00590

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N9/50 A61K38/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HARRACH T ET AL: "Isolation and partial characterization of basic proteinases from stem bromelain." JOURNAL OF PROTEIN CHEMISTRY, vol. 14, no. 1, January 1995, pages 41-52, XP002069063	1-8, 14-19
Y	see the whole document ---	9-19
Y	WO 96 00082 A (CORTECS LTD ;MYNOTT TRACEY LEAHANNE (GB); ENGWERDA CHRISTIAN (GB)) 4 January 1996 see the whole document ---	9-19
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

1 July 1998

Date of mailing of the international search report

13/07/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Van der Schaal, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/00590

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
Y	MUNZIG E ET AL: "Bromelain protease F9 reduces the CD44 mediated adhesion of human peripheral blood lymphocytes to human umbilical vein endothelial cells" FEBS LETTERS, vol. 351, 1994, pages 215-218, XP002069064 see the whole document	9-19
X	ROWAN A ET AL: "Ananain: a novel cysteine proteinase found in pineapple stem" ARCHIVES OF BIOCHEMISTRY AND BIOPHYSICS, vol. 267, no. 1, 15 November 1988, pages 262 -270, XP002069065 see the whole document	1-8
X	EP 0 313 346 A (GENZYME CORP) 26 April 1989 see the whole document	1-8, 16-19

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/00590

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9600082 A	04-01-1996	AU 2749395 A	19-01-1996
		CA 2193654 A	04-01-1996
		EP 0766565 A	09-04-1997
		FI 965204 A	21-02-1997
		JP 10502073 T	24-02-1998
		NO 965564 A	24-02-1997
EP 0313346 A	26-04-1989	AU 628790 B	24-09-1992
		AU 2405988 A	08-06-1989
		CA 1319121 A	15-06-1993
		DE 3888082 D	07-04-1994
		DE 3888082 T	09-06-1994
		DK 589388 A	23-04-1989
		ES 2061682 T	16-12-1994
		JP 2016977 A	19-01-1990
		JP 2644006 B	25-08-1997
		US 5106621 A	21-04-1992

26x2